


SPECIFICATION

Process for Synthesizing 4-Substituted Azetidinone Derivatives


5 Technical Field
[Technical Field]

10 A The present invention relates to a process for producing 4-substituted azetidinone derivatives which are important as a synthetic intermediate for carbapenem based antimicrobial agents and the like.

15 A Background Art
There have already been reported several useful processes (for example, Japanese Unexamined Patent Publication No. 61-207373) for producing azetidinone derivatives represented by the general formula (1):



(wherein OR₁ is a protected hydroxyl group; R₂ is a substituted or unsubstituted alkyl group, a substituted or unsubstituted alkenyl group or a substituted or unsubstituted aromatic group). In making an attempt to produce from such azetidinone derivatives of the general formula (1) substituted azetidinone derivatives represented by the general formula [3]:




(wherein OR₁ is as defined above; CO₂R₃ is an esterified carboxyl group; X and Y are the same or different and represent individually a substituted or unsubstituted alkyl group, a substituted or unsubstituted alkenyl group, a

substituted or unsubstituted aralkyl group, a substituted or
unsubstituted aryl group, a substituted or unsubstituted
alkylthio group, a substituted or unsubstituted alkenylthio
group, a substituted or unsubstituted aralkylthio group, a
5 substituted or unsubstituted arylthio group, a substituted or
unsubstituted alkyloxy group, a substituted or unsubstituted
alkenyloxy group, a substituted or unsubstituted aralkyloxy
group, a substituted or unsubstituted aryloxy group, a
substituted or unsubstituted silyloxy group, a substituted or
10 unsubstituted heterocyclic group, a substituted or
unsubstituted heterocyclic-thio group, a substituted or
unsubstituted heterocyclic-oxy group, [a substituted or
unsubstituted acyl group, a substituted or unsubstituted
ester group, a substituted or unsubstituted thioester group,
15 a substituted or unsubstituted amide group, a substituted or
unsubstituted amino group, a hydrogen atom or a halogen atom,
or are taken together with each other to form a substituted
or unsubstituted cycloalkan-2-on-1-yl group), it has been
necessary to oxidize the sulfide group of the azetidinone
20 A derivative of the general formula [1] to thereby convert the
group to the more readily removable sulfone group or to
substitute the acyloxy group for the sulfide group.

In order to convert the azetidinone derivative of the
general formula [1] to the corresponding sulfone-containing
25 derivative, there is known a method utilizing peroxy acid
(Yoshida et al., Chem. Pharm. Bull., 29, 2899 (1981)), while
a method using a mercury compound (Yoshida et al., Chem.
Pharm. Bull., 29, 2899 (1981)) is known for the conversion of
the same to the acyloxy-containing derivative, but
30 difficulties are encountered in bringing these methods into
commercial practice in terms of hygiene, safety or toxicity
and the like.

A Referring to the conversion of [1] to the acyloxy
derivative, there has recently been reported a useful method
35 involving the use of a copper compound (Japanese
Unexamined Patent Publication No. 3-163057), resulting in
marked improvement in the commercial production of the said
derivative. Nevertheless, this method requires heating in
carrying out the reaction and is yet to be improved. On the

other hand, there have been reported ^α method (Japanese Unexamined Patent Publication No. 3-157365) of directly converting the azetidinone derivative of the general formula [1] to the azetidinone derivative of the general formula [3]:



[3]


(wherein OR₁, CO₂R₃, X and Y are as defined above) and also a method (Ito et al., Tetrahedron, 47, 2801 (1991)) of converting an acyloxyazetidinone derivative to an azetidinone derivative of the general formula [3], which methods have been developed with a specific view to introducing a substituent into the 1-methylene group to enhance the chemical and in vivo stabilities of carbapenem based antimicrobial compounds. However, these methods do not allow the ester compound to be utilized directly as a substitution reagent; before the said reactions are carried out, the former requires the ester compound to be converted to the corresponding diazonium reagent, while the latter necessitates the conversion of the same to the oxazolidone reagent, respectively.

A method of converting the azetidinone derivative of the general formula ⁽¹⁾ directly to the azetidinone derivative of the general formula [3] constitutes a process having one step less as compared with the method of converting the azetidinone derivative of ⁽¹⁾ to the sulfone-containing or acyloxy-containing derivative to thereby effect the intended synthesis, and could consequently offer a useful, advantageous synthetic means, only if the method can be brought to practice by a simple and practical operation. If the method permits the reaction to be carried out at lower temperatures and can furthermore allow a readily available reagent to be used in the introduction of a substituent at the 1-methylene group directly without being converted to another reagent, in addition, substantial improvement could

be achieved in practical aspects. In view of the above, the present inventors conducted investigation into a method permitting the azetidinone derivative of the general formula [1] to be converted directly to the azetidinone derivative of the general formula [3] under mild conditions, and have found out the present invention.


Summary of the Invention
~~[Disclosure of the Invention]~~

The present inventors found that an ester compound represented by the general formula ~~(2)~~ ⁽²⁾ [2]:




~~(2)~~ ⁽²⁾ [2]
~~(2)~~ ⁽²⁾

(wherein CO_2R_3 , X and Y are as defined above) can be treated with a metal base to give the corresponding metal enolate, which can then be reacted with an azetidinone derivative represented by the general formula ~~(1)~~ ⁽¹⁾ [1]:



~~(1)~~ ⁽¹⁾ [1]
~~(1)~~ ⁽¹⁾


(wherein OR_1 and R_2 are as defined above) in the presence of a copper compound to produce a 4-substituted azetidinone derivative represented by the general formula [3]:



[3]

(wherein OR_1 , CO_2R_3 , X and Y are as defined above).

At the same time, it was found that an ester compound represented by the general formula ~~(2)~~ ⁽²⁾ [2]:




(2)
A ~~(2)~~
^

(wherein CO_2R_3 , X and Y are as defined above) can be reacted with an azetidinone derivative represented by the general

formula ~~(1)~~
A ~~(1)~~
^


5



(1)
A ~~(1)~~
^

(wherein OR_1 and R_2 are as defined above) in the presence of zinc and copper compounds to produce a 4-substituted azetidinone derivative represented by the general formula [3]:

15




[3]

(wherein OR_1 , CO_2R_3 , X and Y are as defined above).

With reference to the decarboxylation reaction of 4-substituted azetidinone derivatives represented by the general formula ~~(4)~~
A ~~(4)~~
^

25




(4)
A ~~(4)~~
^

(wherein OR_1 , CO_2R_3 , X and Y are as defined above; R_4 is a hydrogen atom or a protective group or a substituent group for amino group), there has been reported a method (Japanese Unexamined Publication No. 5-155850) involving the

selection of a type of substituents for R₄ to thereby control the configurations of X and Y, and by virtue of the method, highly industrially useful 1-β-substituted derivatives

can be produced preferentially. Consequently, the 4-substituted azetidinone derivatives of the general formula ~~[3]~~⁽³⁾ as synthesized by the reaction according to the present invention and the 4-substituted azetidinone derivatives of the general formula ~~[4]~~⁽⁴⁾ wherein R₄ is a protective group or a substituent group for an amino group readily derived from those by conventional methods, and the 4-substituted azetidinone derivatives of the general formula ~~[5]~~⁽⁵⁾



(wherein OR₁, R₄ and Y are as defined above; X' is the same as defined for X or a mercapto, hydroxyl, formyl, carboxyl or thiocarboxyl group), which are obtained by converting such derivative to a carboxylic acid compound, followed by decarboxylation, can be utilized as a starting compound for carbapenem compounds of utility as a therapeutic agent for infections.

Such being the case, it was also found that the 4-substituted azetidinone derivatives of the general formulae [3] and [4] are converted to their carboxylic acid compounds for example by way of hydrolysis or a reaction procedure selected for the type of esters, followed by decarboxylation treatment to give the 4-substituted azetidinone derivatives of the general formula ~~[5]~~⁽⁵⁾ useful as a starting material for carbapenem compounds.

The present invention is based on such novel findings and relates to (A) a process for synthesizing 4-substituted azetidinone derivatives of the above-described general formula ~~[3]~~⁽³⁾, characterized in that said process comprises reacting an azetidinone derivative of the above-described general formula ~~[1]~~⁽¹⁾ with an ester compound of the above-

A described general formula ⁽²⁾[2] in the presence of zinc and copper compounds, (B) a process for synthesizing 4-substituted azetidinone derivatives of the above-described general formula [3], characterized in that said process comprises
 5 treating an ester compound of the above-described general formula ⁽²⁾[2] with a metal base to give a metal enolate, which is then reacted with an azetidinone derivative of the above-mentioned general formula ⁽¹⁾[1] in the presence of a copper compound, (C) a process for synthesizing 4-substituted
 10 azetidinone derivatives of the above-mentioned general formula ⁽³⁾[3], characterized in that said process comprises converting an ester compound of the above-mentioned general formula ⁽³⁾[3] and ⁽⁴⁾[4] to a carboxylic acid compound, followed by decarboxylation treatment, and (D) novel 4-substituted
 15 azetidinone derivatives among the compounds represented by the above-mentioned general formulae ⁽³⁾[3], ⁽⁴⁾[4] and ⁽⁵⁾[5] in the above-mentioned processes.

The substitution reaction for the 4-position of the
 20 A azetidinone derivative of the general formula ⁽¹⁾[1] is normally carried out in hydrocarbon solvents such as benzene and toluene, chlorinated hydrocarbon solvents such as methylene chloride and chloroform, nitrile solvents such as acetonitrile, ketone solvents such as acetone and methyl vinyl ketone, ether solvents such as diethyl ether and
 25 tetrahydrofuran, and ester solvents such as ethyl acetate, either solely or in mixtures thereof.

A The ester compound of the general formula ⁽²⁾[2] is preferably used in proportions of 1 to 3 equivalents against
 30 A the azetidinone derivative of the general formula ⁽¹⁾[1]. The ester compound is preferably acted on by an equal equivalent of a metal base to produce a metal enolate, followed by addition of the azetidinone derivative of the general formula [1] and a copper compound to allow the reaction.

Alternatively, when the ester compound of the general formula
 35 A ⁽²⁾[2] is reacted with the azetidinone derivative of the general formula ⁽¹⁾[1] directly without converting to the corresponding metal enolate, the reaction is effected in the above-mentioned solvents, wherein such reaction can be conducted in the presence of zinc and copper compounds by ⁽²⁾preferably using

the ester compound of the general formula [2] in proportions of 1 to 3 equivalents against the azetidinone derivative of the general formula [1].

As the copper compound, there may be mentioned, for example, copper oxides, copper halides, salts of copper with organic carboxylic acids, salts of copper with mineral acids, their complexes and the like. Their preferred examples include cuprous oxide, cupric oxide, cuprous chloride, cupric chloride, cuprous bromide, cupric fluoride, cuprous iodide, cupric perchlorate, cupric nitrate, cupric sulfate, cuprous sulfide, cupric sulfide, cupric trifluoromethanesulfonate, copper cyanide, salts of copper with aliphatic carboxylic acids such as cuprous acetate, cupric acetate, cupric trifluoroacetate, copper propionate and copper butyrate, salts of copper with aromatic carboxylic acids such as copper benzoate, etc., and there are usually employed cuprous halides and their complexes, with cuprous bromide dimethyl sulfide complex being particularly preferred. The used amount each of the copper compounds and zinc is suitably in the range of 1 to 4 equivalents against the azetidinone derivative of the general formula [1]. Meanwhile, preferred examples of the metal base include alkali metal hydride compounds such as sodium hydride and potassium hydride. The reaction temperature varies depending upon the type of the used ester compound of the general formula [2], kind of the azetidinone derivative of the general formula [1], and the like, and is usually in the range of 0°C to 50°C, suitably in the region of 0°C to room temperature.

The conversion of the 4-substituted azetidinone derivative of the general formula ~~(3)~~ to the 4-substituted azetidinone derivative of the general formula ~~(4)~~ where R₄ is a protective group or a substituent group for the amino group can be done by the conventional procedures.

Referring to the post-reaction treatment, to the reaction solution is added a saturated ammonium chloride solution, followed by extraction of the solution mixture with an organic solvent, and the organic extract layer can be washed with water, dried and concentrated to give the objective compound. The crude reaction product can be used in

the subsequent reaction directly without being purified, but can also be purified by recrystallization or column chromatography, if necessary.

5 The conversion reaction of the ester compound of the
general formulae ⁽³⁾~~[3]~~ and ⁽⁴⁾~~[4]~~ to the corresponding carboxylic
acid compound is usually carried out, in pyridine based
solvents such as pyridine, lutidine and collidine, nitrile
based solvents such as acetonitrile, ketone based solvents
such as acetone and methyl vinyl ketone, ether based solvents
10 such as tetrahydrofuran, and alcohol based solvents such as
methanol and ethanol, either solely or in solvent mixtures
thereof, suitably in pyridine, collidine or tetrahydrofuran,
by hydrolysis in the presence of a base, for example, by
adding an aqueous solution of a metal hydroxide compound to
15 the compound of ~~the~~ general formulae ⁽³⁾~~[3]~~ and ⁽⁴⁾~~[4]~~ or under the
conditions of a selective procedure adopted according to
the type of the ester compound.

The conditions of the selective reaction procedure
according to the type of the ester compounds can be
20 exemplified by the reaction utilizing metals such as
palladium for allyl ester, the catalytic hydrogenation
procedures for benzyl ester, the procedures utilizing zinc
for trichloroethyl ester, and the like, and these all are
known and can be carried out in accordance with the
25 descriptions given in various pieces of literature and the
like.

The metal hydroxide compound is preferably used in
proportions of 1 to 2 equivalents against the ester compound
to conduct the reaction at a temperature in the range of
30 normally 0°C to 50°C, preferably 0°C to room temperature.
Examples of the metal hydroxide compound include alkali metal
hydroxide compounds such as sodium hydroxide, potassium
hydroxide and lithium hydroxide, and alkaline earth metal
hydroxide compounds such as barium hydroxide.

35 The decarboxylation reaction of the carboxylic acid
compound to form the 4-substituted azetidinone derivative of
the general formula ⁽⁵⁾~~[5]~~ can be carried out, for example, by
adding an acid in an amount equivalent to the metal hydroxide
compound used in the hydrolysis to give the carboxylic acid

compound, followed by heating normally at 50°C to 200°C, preferably 100°C to 150°C. The acid can be exemplified by mineral acids such as hydrochloric acid, sulfuric acid and nitric acid, organic acids such as acetic acid, propionic acid and camphorsulfonic acid, an aqueous solution thereof, ion exchange resins, etc.

The post-reaction treatment can be effected by concentrating the reaction solution, extracting the concentrate with an organic solvent and washing the organic extract layer with water, followed by drying and concentrating to produce the objective compound. The crude reaction product can be used in the subsequent reaction directly without being purified, but can also be purified by recrystallization or column chromatography, if necessary.

The removal procedure of the protective group for the amino group in R₄ of the 4-substituted azetidinone derivative of the general formula (3) varies with the type of the protective group, and can be brought to practice by selecting suitably an appropriate reaction; when the protective group is a tri-substituted silyl group, for example, a weak acid such as dilute hydrochloric acid may be reacted with the azetidinone derivative of (5), while in the case of the protective group being benzyl, phenethyl or benzhydryl group which may be substituted, metallic sodium may be reacted with the derivative of (5) in liquid ammonia by the Barch's reduction.

The 4-substituted azetidinone derivative of the general formula (5) as produced according to the present invention can be utilized, directly or after removal or replacement of the protective group for the amino group in R₄, as a starting compound for carbapenem based compounds.

The conversion procedure to a carbapenem based compound varies depending upon the kind of substituents at the 4-position, and can be conducted into practice by a suitably selected reaction, for example, in accordance with the descriptions on the cyclization reaction (Hatanaka, M. et al., Tetrahedron Lett., 1981, 22, 3883) by way of Diechmann reaction, reductive cyclization reaction (Shibata, T. et al., J. Antibiotics, 1989, 42, 374) with use of phosphite,

cyclization reaction (Guthikonda, R. N. et al., J. Med. Chem., 1987, 30, 871) based on Wittig reaction, cyclization reaction with use of transitional metals (Ratcliffe, R. W. et al., Tetrahedron Lett., 1980, 21, 1193) and the like.

- 5 The protected hydroxyl group represented by OR₁ is not particularly limited and comprehends hydroxyl groups protected with protective groups conventionally used for the hydroxyl group, which are exemplified by tri-substituted silyloxy groups, specifically trialkylsilyloxy, 10 aryl(alkyl)alkoxysilyloxy, alkoxydiarylsilyloxy, triarylsilyloxy, alkyl-diarylsilyloxy, aryldialkylsilyloxy, and triaralkylsilyloxy groups, etc., such as trimethylsilyloxy, triethylsilyloxy, triisopropylsilyloxy, dimethylhexylsilyloxy, tert-butyl-dimethylsilyloxy, 15 methyldiisopropylsilyloxy, isopropyl-dimethylsilyloxy, tert-butylmethoxyphenylsilyloxy, tert-butoxydiphenylsilyloxy, triphenylsilyloxy, tert-butyl-diphenylsilyloxy, dimethylcumylsilyloxy and tribenzylsilyloxy groups, etc.; lower alkoxy groups which may have at least one suitable 20 substituent, such as methoxymethoxy, methoxyethoxymethoxy and triphenylmethoxy groups; lower alkanoyloxy groups which may have at least one suitable substituent, such as acetoxy, chloroacetoxy, methoxyacetoxy, propionyloxy, butyryloxy, isobutyryloxy, valeryloxy, pivaloyloxy, hexanoyloxy, 2- 25 ethylbutyryloxy, 3,3-dimethylbutyryloxy and pentanoyloxy groups; lower alkoxycarbonyloxy groups which may have at least one suitable substituent, such as methoxycarbonyloxy, ethoxycarbonyloxy, propoxycarbonyloxy, isopropoxycarbonyloxy, tert-butoxycarbonyloxy, 2-iodoethoxycarbonyloxy, 2,2- 30 dichloroethoxycarbonyloxy and 2,2,2-trichloroethoxycarbonyloxy groups; lower alkenyloxycarbonyloxy groups which may have at least one suitable substituent, such as vinyloxycarbonyloxy, allyloxycarbonyloxy and 2-chloroallyloxycarbonyloxy groups; arylcarbonyloxy groups which may have at least one 35 suitable substituent, such as benzoyloxy group;
A aralkyloxycarbonyloxy ^{alkyl} groups which may have at least one suitable substituent, such as benzyloxycarbonyloxy, p-nitrobenzyloxycarbonyloxy, p-methoxybenzyloxycarbonyloxy, phenethyloxycarbonyloxy, trityloxycarbonyloxy, benzhydryl-

oxycarbonyloxy, bis(methoxyphenyl)methyloxycarbonyloxy, 3,4-dimethoxybenzyloxycarbonyloxy and 4-hydroxy-3,5-di-tert-butylbenzyloxycarbonyloxy groups; aryloxycarbonyloxy groups which may have at least one suitable substituent, such as phenyloxycarbonyloxy, 4-chlorophenyloxycabonyloxy, tolyloxycarbonyloxy, tert-butylphenyloxycarbonyloxy, xylyloxycarbonyloxy, mesityloxycarbonyloxy and cumenyloxy-carbonyloxy groups; aralkyloxy groups which may have at least one suitable substituent, such as benzyloxy, p-nitrobenzyloxy, p-methoxybenzyloxy, p-tert-butylbenzyloxy, 3,4-dimethylbenzyloxy, 2,4-dimethoxybenzyloxy, benzhydryloxy and trityloxy groups; heterocyclic-oxy groups which may have at least one suitable substituent, such as tetrahydropyranyloxy group. Unless otherwise specified particularly, the term "lower" designates the same meaning in the below description throughout the present specification, and means preferably 1 to 6 in number of carbon atoms, particularly preferably 1 to 4 in number of carbon atoms.

The group represented by R₂ is not particularly limited, but examples of such group, which are readily available and can be produced at low costs, include alkyl groups being exemplified by linear or branched lower alkyl groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, tert-butyl and hexyl groups and monocyclic or polycyclic alkyl groups such as cyclopentyl, cyclohexyl, menthyl, fenchyl and bornyl groups; alkenyl groups being exemplified by linear or branched lower alkenyl groups such as vinyl, allyl, 1-propenyl, 2-butenyl and 2-methyl-2-propenyl groups; aromatic groups being exemplified by aryl groups having a number of carbon atoms of 6 to 10 such as phenyl, tolyl, xylyl, mesityl and cumenyl groups and aromatic heterocyclic groups such as pyridyl, pyrimidinyl, pyrazinyl and pyridazinyl groups, wherein such alkyl, alkenyl, and aromatic groups individually may be substituted with one or not less than two substituents, for example, halogen atoms such as fluorine, chlorine and bromine atoms; linear or branched lower alkyl groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, tert-butyl and hexyl groups; monocyclic or polycyclic alkyl groups such as cyclopentyl, cyclohexyl, menthyl, fenchyl and

bornyl groups; linear or branched lower alkoxy groups such as methoxy and ethoxy groups; carboxyl group; amino group; nitro group; cyano group; hydroxyl group; aryl groups having a number of carbon atoms of 6 to 10, such as phenyl, tolyl, xylyl, mesityl and cumenyl groups, which may be substituted with the above-mentioned halogen atoms, and lower alkyl, lower alkoxy, carboxyl, amino, nitro, cyano and hydroxyl groups; aralkyl groups having a number of carbon atoms of 7 to 24, such as benzyl, phenethyl, trityl and benzhydryl groups, which may be substituted with the above-mentioned halogen atoms, and lower alkyl, lower alkoxy, carboxyl, amino, nitro, cyano and hydroxyl groups.

The group represented by R_3 is not particularly limited, only if it can eliminate from the esterified carboxyl group represented by CO_2R_3 through hydrolysis or under conditions of the selective procedure according to the type of esters, and its preferred examples include those capable of forming the following esters:

Tri-substituted silyl esters such as trialkylsilyl esters, aryl(alkyl)alkoxysilyl esters, alkoxydiarylsilyl esters, triarylsilyl esters, alkyl diarylsilyl esters, aryldialkylsilyl esters, triaralkylsilyl esters (e.g. trimethylsilyl, triethylsilyl, triisopropylsilyl, dimethylhexylsilyl, tert-butyl dimethylsilyl, methyl-diisopropylsilyl, isopropyl dimethylsilyl, tert-butyl-methoxyphenylsilyl, tert-butoxydiphenylsilyl, triphenylsilyl, tert-butyl diphenylsilyl, dimethylcumylsilyl and tribenzylsilyl esters); tri-substituted silyl lower-alkyl esters such as trialkylsilyl lower-alkyl esters, aryl(alkyl)alkoxysilyl lower-alkyl esters, alkoxydiarylsilyl lower-alkyl esters, triarylsilyl lower-alkyl esters, alkyl diarylsilyl lower-alkyl esters, aryldialkylsilyl lower-alkyl esters, and triaralkylsilyl lower-alkyl esters (e.g. the above exemplified compounds in which the tri-substituted silyl groups are substituted with lower alkyl groups (e.g. linear or branched lower alkyl such as methyl, ethyl, n-propyl, isopropyl, n-butyl, tert-butyl, ^{or hexyl} aromatic heterocyclic esters (e.g. pyridyl, pyrimidinyl, pyrazinyl and pyridazinyl esters); lower alkyl esters (e.g. methyl,

ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl and hexyl esters), lower alkyl esters which may have at least one suitable substituent, such as lower alkanoyloxy (lower)alkyl esters [e.g. acetoxymethyl, propionyloxymethyl, butyryloxymethyl, valeryloxymethyl, pivaloyloxymethyl, hexanoyloxymethyl, 1-(or 2-)acetoxylethyl, 1- (or 2- or -3)acetoxypentyl, 1-(or 2- or 3- or 4-)acetoxylbutyl, 1-(or 2-)propionyloxyethyl, 1-(or 2- or 3-)propionyloxypropyl, 1-(or 2-)butyryloxyethyl, 1- (or 2-)isobutyryloxyethyl, 1- (or 2-)pivaloyloxyethyl, 1-(or 2-)hexanoyloxyethyl, isobutyryloxymethyl, 2-ethylbutyryloxymethyl, 3,3-dimethylbutyryloxymethyl and 1-(or 2-)pentanoyloxyethyl esters], lower alkanesulfonyl (lower)alkyl esters (e.g. 2-mesyloethyl ester), mono (or di or tri)halo(lower)alkyl esters (e.g. 2-iodoethyl, 2,2-dichloroethyl and 2,2,2-trichloroethyl esters), lower alkoxy-carbonyloxy(lower)alkyl esters [e.g. methoxycarbonyloxymethyl, ethoxycarbonyloxymethyl, propoxycarbonyloxymethyl, tert-butoxycarbonyloxymethyl, 1- (or 2-)methoxycarbonyloxyethyl, 1- (or 2-) ethoxycarbonyloxyethyl and 1- (or 2-)isopropoxycarbonyloxyethyl esters]. phthalidylidene (lower)alkyl esters, or (5-lower-alkyl-2-oxo-1,3-dioxolene-4-yl)(lower)alkyl esters [e.g. (5-methyl-2-oxo-1,3-dioxolene-4-yl)methyl, (5-ethyl-2-oxo-1,3-dioxolene-4-yl)methyl and (5-propyl-2-oxo-1,3-dioxolene-4-yl) ethyl esters]; lower alkenyl esters (e.g. vinyl and allyl esters); lower alkynyl esters (e.g. ethynyl and propynyl esters); aryl(lower)alkyl esters which may have at least one suitable substituent [e.g. benzyl, 4-methoxybenzyl, 4-nitrobenzyl, phenethyl, trityl, benzhydryl, bis(methoxyphenyl)methyl, 3,4-dimethoxybenzyl and 4-hydroxy-3,5-di-tert-butylbenzyl esters]; aryl esters which may have at least one suitable substituent (e.g. phenyl, 4-chlorophenyl, tolyl, tert-butylphenyl, xylyl, mesityl and cumenyl esters); phthalidyl esters, etc.

Furthermore, these may be substituted at their individual groups with one or more substituents, being exemplified by halogen atoms such as fluorine, chlorine and bromine atoms; linear or branched lower alkyl groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, tert-butyl and

hexyl groups; monocyclic or polycyclic alkyl groups such as cyclopentyl, cyclohexyl, menthyl, fenchyl and bornyl groups; linear or branched lower alkoxy groups such as methoxy and ethoxy groups; carboxyl group; amino group; nitro group; cyano group; hydroxyl group; aryl groups having a number of carbon atoms of 6 to 10 which may be substituted with the above-mentioned halogen atoms and lower alkyl, lower alkoxy, carboxyl, amino, nitro, cyano and hydroxyl groups, etc., such as phenyl, tolyl, xylyl, mesityl and cumenyl; aralkyl groups having a number of carbon atoms of 7 to 24 which may be substituted with the above-mentioned halogen atoms and lower alkyl, lower alkoxy, carboxyl, amino, nitro, cyano and hydroxyl groups, such as benzyl, phenethyl, trityl and benzhydryl groups; and the below-described heterocyclic groups, acyl and ester groups.

Preferred examples of the groups represented by X and Y include the groups which can be adopted as a synthetic intermediate for carbapenem based antimicrobial compounds, being specifically exemplified by hydrogen atoms and the below-described groups:

Referring to the alkyl, alkenyl, aralkyl, aryl, alkylthio, alkenylthio, aralkylthio, arylthio, alkyloxy, alkenyloxy, aralkyloxy and aryloxy groups, namely, as the alkyl group, there are mentioned linear or branched lower alkyl groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, tert-butyl and hexyl groups, and monocyclic or polycyclic alkyl groups such as cyclopentyl, cyclohexyl, menthyl, fenchyl and bornyl groups; the alkenyl group includes for example linear or branched lower alkenyl groups such as vinyl, allyl, 1-propenyl, 2-butenyl and 2-methyl-2-propenyl groups; examples of the aralkyl group include aralkyl groups having a number of carbon atoms of 7 to 24 such as benzyl, phenethyl, trityl and benzhydryl groups; and as the aryl group, there may be mentioned aryl groups having a number of carbon atoms of 6 to 10 such as phenyl, tolyl, xylyl, mesityl and cumenyl groups.

As the silyloxy group, there may be mentioned tri-substituted silyloxy groups, and their specific examples include trialkylsilyloxy, aryl(alkyl)alkoxysilyloxy,

- alkoxydiarylsilyloxy, triarylsilyloxy, alkyl-diarylsilyloxy, aryl-dialkylsilyloxy and triaralkylsilyloxy groups, being exemplified by trimethylsilyloxy, triethylsilyloxy, triisopropylsilyloxy, dimethylhexylsilyloxy, tert-butyl-dimethylsilyloxy, methyl-diisopropylsilyloxy, isopropyl-dimethylsilyloxy, tert-butylmethoxyphenylsilyloxy, tert-butoxydiphenylsilyloxy, triphenylsilyloxy, tert-butyl-diphenylsilyloxy, dimethylcumylsilyloxy and tribenzylsilyloxy groups.
- These alkyl, alkenyl, aralkyl, aryl, alkylthio, alkenylthio, aralkylthio, arylthio, alkyloxy, alkenyloxy, aralkyloxy, aryloxy and silyloxy groups may be substituted in their individual groups with one or more substituents, for example, halogen atoms such as fluorine, chlorine and bromine atoms; carboxyl group; formyl group; nitro group; cyano group; hydroxyl group; amino group; linear or branched lower alkyl groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, tert-butyl and hexyl groups; monocyclic and polycyclic alkyl groups such as cyclopentyl, cyclohexyl, menthyl, fenchyl and bornyl groups; linear or branched lower alkenyl groups such as vinyl, allyl, 2-chloroallyl, 1-propenyl, 2-butenyl, and 2-methyl-2-propenyl groups; aryl groups having a number of carbon atoms of 6 to 10 such as phenyl, tolyl, xylyl, mesityl and cumenyl groups; aralkyl groups having a number of carbon atoms of 7 to 24 such as benzyl, phenethyl, trityl and benzhydryl groups; alkylthio, alkenylthio, aralkylthio, arylthio, alkyloxy, alkenyloxy, aralkyloxy and aryloxy groups which correspond individually to the above-mentioned alkyl, alkenyl, aralkyl and aryl groups;
- alkylsulfinyl and alkylsulfonyl groups that correspond individually to the above-mentioned alkyl groups; aralkylsulfinyl and aralkylsulfonyl groups that correspond individually to the above-mentioned aralkyl groups; arylsulfinyl and arylsulfonyl groups that correspond individually to the above-mentioned aryl groups; carbamoyl groups; carbamoyloxy groups; imino-lower-alkyl groups; imino-lower-alkylamino groups; acyloxy groups that correspond individually to the below-described acyl groups; the above-mentioned silyloxy groups; the below-described heterocyclic,

heterocyclic-thio, heterocyclic-oxy, acyl, ester, thioester and amide groups.

Furthermore, the above-described substituents individually may be substituted with one or more of substituents such as the above-described substituents: by way of examples, the substituents for the alkyl groups (as is the same with the alkylthio and alkyloxy groups) include halogen atoms, and carboxyl, formyl, nitro, cyano, hydroxyl, amino, alkyl, alkenyl, aryl, aralkyl, alkylthio, alkenylthio, aralkylthio, arylthio, alkyloxy, alkenyloxy, aralkyloxy, aryloxy, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, carbamoyl, carbamoyloxy, imino, imino-lower-alkylamino, acyloxy, silyloxy, heterocyclic, heterocyclic-thio, heterocyclic-oxy, acyl, ester, thioester and amide groups; as the substituents for the alkenyl groups (as is the same with the alkenylthio and alkenyloxy groups), there may be mentioned halogen atoms, and carboxyl, formyl, nitro, cyano, hydroxyl, amino, alkyl, aryl, aralkyl, alkylthio, alkenylthio, aralkylthio, arylthio, alkyloxy, alkenyloxy, aralkyloxy, aryloxy, alkylsulfinyl, alkyl-sulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, carbamoyl, carbamoyloxy, imino-lower-alkyl, imino-lower-alkylamino, acyloxy, silyloxy, heterocyclic, heterocyclic-thio, heterocyclic-oxy, acyl, ester, thioester and amide groups; the substituents for the aralkyl groups (as is the same with the aralkylthio and aralkyloxy groups) include for example halogen atoms, and carboxyl, formyl, nitro, cyano, hydroxyl, amino, alkyl, alkenyl, aryl, aralkyl, alkylthio, alkenylthio, aralkylthio, arylthio, alkyloxy, alkenyloxy, aralkyloxy, aryloxy, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, carbamoyl, carbamoyloxy, imino-lower-alkyl, imino-lower-alkylamino, acyloxy, silyloxy, heterocyclic, heterocyclic-thio, heterocyclic-oxy, acyl, ester, thioester and amide groups; as the substituents for the aryl groups (as is the same with the arylthio and aryloxy groups), there may be mentioned halogen atoms, and carboxyl, formyl, nitro, cyano, hydroxyl, amino, alkyl, alkenyl, aryl, aralkyl, alkylthio, alkenylthio,

aralkylthio, arylthio, alkyloxy, alkenyloxy, aralkyloxy,
aryloxy, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl,
aralkylsulfonyl, arylsulfinyl, arylsulfonyl, carbamoyl,
carbamoyloxy, imino-lower-alkyl, imino-lower-alkylamino,
5 acyloxy, silyloxy, heterocyclic, heterocyclic-thio,
heterocyclic-oxy, acyl, ester, thioester and amide groups;
and the substituents for the amino, carbamoyl, carbamoyloxy,
imino-lower-alkyl, imino-lower-alkylamino and amide groups
include for example halogen atoms, and carboxyl, formyl,
10 nitro, cyano, hydroxyl, amino, alkyl, alkenyl, aryl, aralkyl,
alkylthio, alkenylthio, aralkylthio, arylthio, alkyloxy,
alkenyloxy, aralkyloxy, aryloxy, alkylsulfinyl,
alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl,
arylsulfinyl, arylsulfonyl, carbamoyl, carbamoyloxy, imino-
15 lower-alkyl, imino-lower-alkylamino, acyloxy, silyloxy,
heterocyclic, heterocyclic-thio, heterocyclic-oxy, acyl,
ester, thioester and amide groups.

The term "heterocyclic group" in the heterocyclic,
heterocyclic-thio and heterocyclic-oxy groups is understood
20 to comprehend saturated or unsaturated, monocyclic or
polycyclic heterocyclic groups having at least one hetero
atom such as oxygen, sulfur and nitrogen atoms, and their
preferred examples include 3- to 8-membered, particularly
preferably 5- or 6-membered unsaturated monocyclic
25 heterocyclic groups having 1 to 4 nitrogen atoms such as
pyrrolyl, pyrrolinyl, imidazolyl, pyrazolyl and pyridyl
groups and their N-oxides, pyrimidyl, pyrazinyl, pyridazinyl,
triazolyl (e.g. 4H-1,2,4-triazolyl, 1H-1,2,3-triazolyl and
2H-1,2,3-triazolyl groups), tetrazolyl (e.g. 1H-tetrazolyl
30 and 2H-tetrazolyl groups) and dihydrotriazinyl (e.g. 4,5-
dihydro-1,2,4-triazinyl and 2,5-dihydro-1,2,4-triazinyl
groups) groups; 3- to 8-membered, particularly preferably 5-
or 6-membered saturated monocyclic heterocyclic groups having
1 to 4 nitrogen atoms such as azetidiny, pyrrolidinyl,
35 imidazolidinyl, piperizinyl, pyrazolidinyl and piperazinyl
groups; 7- to 12-membered, unsaturated polycyclic
heterocyclic groups having 1 to 5 nitrogen atoms such as
indolyl, isoindolyl, indolizinyl, benzimidazolyl, quinolyl,
isoquinolyl, indazolyl, benzotriazolyl, tetrazopyridyl,

- 5 tetrazopyridazinyl (e.g. tetrazo[1,5-b]pyridazinyl group)
and dihydrotriazolopyridazinyl; 3- to 8-membered,
particularly preferably 5- or 6-membered unsaturated
monocyclic heterocyclic groups having 1 to 2 oxygen atoms and
1 to 3 nitrogen atoms such as oxazolyl, isoxazolyl and
oxadiazolyl (e.g. 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl and
1,2,5-oxadiazolyl groups) groups; 3- to 8-membered,
particularly preferably 5- or 6-membered saturated monocyclic
heterocyclic groups having 1 to 2 oxygen atoms and 1 to 3
10 nitrogen atoms such as morpholinyl group; 7- to 12-membered
unsaturated polycyclic heterocyclic groups having 1 to 2
oxygen atoms and 1 to 3 nitrogen atoms such as benzoxazolyl
and benzoxadiazolyl groups; 3- to 8-membered, particularly
preferably 5- or 6-membered unsaturated monocyclic
15 heterocyclic groups having 1 to 2 sulfur atoms and 1 to 3
nitrogen atoms such as 1,3-thiazolyl, 1,2-thiazolyl,
thiazolinyl and thiadiazolyl (e.g. 1,2,4-thiadiazolyl,
1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl and 1,2,3-thiadiazolyl
groups); 3- to 8-membered saturated monocyclic heterocyclic
20 groups having 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms
such as thiazolidinyl group; 7- to 12-membered unsaturated
polycyclic heterocyclic groups having 1 to 2 sulfur atoms and
1 to 3 nitrogen atoms such as benzothiazolyl and
benzothiadiazolyl groups; 3- to 8-membered, particularly
25 preferably 5- or 6-membered unsaturated monocyclic
heterocyclic groups having 1 to 2 oxygen atoms such as
furanyl and pyranyl groups; 3- to 8-membered, particularly
preferably 5- or 6-membered saturated monocyclic heterocyclic
groups having 1 to 2 oxygen atoms such as tetrahydrofuranyl
30 and tetrahydropyranyl groups; 3- to 8-membered, particularly
preferably 5- or 6-membered unsaturated monocyclic
heterocyclic groups having a sulfur atom such as thienyl
group and S-oxide; and 3- to 8-membered, particularly
preferably 5- or 6-membered saturated monocyclic heterocyclic
35 groups having a sulfur atom such as tetrahydrothienyl group
and S-oxide.

These heterocyclic groups may be substituted in their
individual groups with one or more substituents, for example,
halogen atoms such as fluorine, chlorine and bromine atoms;


carboxyl group; formyl group; nitro group; cyano group;
hydroxyl group; amino group; linear or branched lower alkyl
groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl,
tert-butyl and hexyl groups; monocyclic and polycyclic alkyl
5 groups such as cyclopentyl, cyclohexyl, menthyl, fenchyl and
bornyl groups; linear or branched lower alkenyl groups such
as vinyl, allyl, 2-chloroallyl, 1-propenyl, 2-butenyl, and 2-
methyl-2-propenyl groups; aryl groups having a number of
10 carbon atoms of 6 to 10 such as phenyl, tolyl, xylyl, mesityl
and cumenyl groups; aralkyl groups having a number of carbon
atoms of 7 to 24 such as benzyl, phenethyl, trityl and
benzhydryl groups; alkylthio, alkenylthio, aralkylthio,
arylthio, alkyloxy, alkenyloxy, aralkyloxy and aryloxy groups
which correspond individually to the above-mentioned alkyl,
15 alkenyl, aralkyl and aryl groups; alkylsulfinyl and
alkylsulfonyl groups that correspond individually to the
above-mentioned alkyl groups; aralkylsulfinyl and
aralkylsulfonyl groups that correspond individually to the
above-mentioned aralkyl groups; arylsulfinyl and arylsulfonyl
20 groups that correspond individually to the above-mentioned
aryl groups; carbamoyl groups; carbamoyloxy groups; imino-
lower-alkyl groups; imino-lower-alkylamino groups; acyloxy
groups that correspond individually to the below-described
acyl groups; the above-mentioned silyloxy, heterocyclic,
25 heterocyclic-thio, heterocyclic-oxy groups; the below-
described acyl, ester, thioester and amide groups.

Furthermore, the above-described substituents
individually may be substituted with one or more of
substituents such as the above-described substituents: by way
30 of examples, the substituents for the alkyl groups (as is the
same with the alkylthio and alkyloxy groups) include halogen
atoms, and carboxyl, formyl, nitro, cyano, hydroxyl, amino,
alkyl, alkenyl, aryl, aralkyl, alkylthio, alkenylthio,
aralkylthio, arylthio, alkyloxy, alkenyloxy, aralkyloxy,
35 aryloxy, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl,
aralkylsulfonyl, arylsulfinyl, arylsulfonyl, carbamoyl,
carbamoyloxy, imino, imino-lower-alkylamino, acyloxy,
silyloxy, heterocyclic, heterocyclic-thio, heterocyclic-oxy,
acyl, ester, thioester and amide groups; as the substituents

for the alkenyl groups (as is the same with the alkenylthio and alkenyloxy groups), there may be mentioned halogen atoms, and carboxyl, formyl, nitro, cyano, hydroxyl, amino, alkyl, aryl, aralkyl, alkylthio, alkenylthio, aralkylthio, arylthio, 5 alkyloxy, alkenyloxy, aralkyloxy, aryloxy, alkylsulfinyl, alkyl-sulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, carbamoyl, carbamoyloxy, imino-lower-alkyl, imino-lower-alkylamino, acyloxy, silyloxy, heterocyclic, heterocyclic-thio, heterocyclic-oxy, acyl, 10 ester, thioester and amide groups; the substituents for the aralkyl groups (as is the same with the aralkylthio and aralkyloxy groups) include for example halogen atoms, and carboxyl, formyl, nitro, cyano, hydroxyl, amino, alkyl, alkenyl, aryl, aralkyl, alkylthio, alkenylthio, aralkylthio, 15 arylthio, alkyloxy, alkenyloxy, aralkyloxy, aryloxy, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, carbamoyl, carbamoyloxy, imino-lower-alkyl, imino-lower-alkylamino, acyloxy, silyloxy, heterocyclic, heterocyclic-thio, 20 heterocyclic-oxy, acyl, ester, thioester and amide groups; as the substituents for the aryl groups (as is the same with the arylthio and aryloxy groups), there may be mentioned halogen atoms, and carboxyl, formyl, nitro, cyano, hydroxyl, amino, alkyl, alkenyl, aryl, aralkyl, alkylthio, alkenylthio, 25 aralkylthio, arylthio, alkyloxy, alkenyloxy, aralkyloxy, aryloxy, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, carbamoyl, carbamoyloxy, imino-lower-alkyl, imino-lower-alkylamino, acyloxy, silyloxy, heterocyclic, heterocyclic-thio, 30 heterocyclic-oxy, acyl, ester, thioester and amide groups; and the substituents for the amino, carbamoyl, carbamoyloxy, imino-lower-alkyl, imino-lower-alkylamino and amide groups include for example halogen atoms, and carboxyl, formyl, nitro, cyano, hydroxyl, amino, alkyl, alkenyl, aryl, aralkyl, 35 alkylthio, alkenylthio, aralkylthio, arylthio, alkyloxy, alkenyloxy, aralkyloxy, aryloxy, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, carbamoyl, carbamoyloxy, imino-lower-alkyl, imino-lower-alkylamino, acyloxy, silyloxy,

heterocyclic, heterocyclic-thio, heterocyclic-oxy, acyl, ester, thioester and amide groups

More specific examples include:



and acyl groups containing them (which have carbonyl group attached to the linkages).

As the acyl group, there may be mentioned alkyl-carbonyl, alkenylcarbonyl, aralkylcarbonyl, arylcarbonyl, alkylthiocarbonyl, alkenylthiocarbonyl, aralkylthiocarbonyl, arylthiocarbonyl, alkylloxycarbonyl, alkenyloxycarbonyl,

aralkyloxycarbonyl, aryloxycarbonyl, silyloxycarbonyl, heterocyclic-carbonyl, heterocyclic-thiocarbonyl, heterocyclic-oxycarbonyl, estercarbonyl, thioestercarbonyl and amidocarbonyl groups which correspond individually to the above-mentioned alkyl, alkenyl, aralkyl, aryl, alkylthio, alkenylthio, aralkylthio, arylthio, alkyloxy, alkenyloxy, aralkyloxy, aryloxy, silyloxy, heterocyclic, heterocyclic-thio and heterocyclic-oxy groups, as well as the below-described ester, thioester and amide groups.

As the ester and thioester groups, furthermore, there may be mentioned carboxyl and thiocarboxyl groups which are esterified with the above-mentioned alkyl, alkenyl, aralkyl, aryl, alkylthio, alkenylthio, aralkylthio, arylthio, alkyloxy, alkenyloxy, aralkyloxy, aryloxy, silyl, heterocyclic, heterocyclic-thio and heterocyclic-oxy or acyl groups, as well as the below-described amide and amino groups, while examples of the amide and amino groups include amide and amino groups which may be substituted individually for example with one or more of the above-mentioned substituents for the amide and amino groups, etc. As the halogen atom, there may be mentioned fluorine, chlorine and bromine atoms.

Referring to the cycloalkan-2-on-1-yl groups which X and Y are taken together with each other to represent, examples of such cycloalkane include monocyclic ones such as cyclopentane and cyclohexane, which may be substituted individually for example with one or more of the above-mentioned substituents for the alkyl groups.

As the groups represented by X', there may be mentioned the groups represented by X as well as mercapto, hydroxyl, formyl, carboxyl and thiocarboxyl groups.

The groups represented by R₄ are not particularly limited, only if they can be adopted as a synthetic intermediate for the carbapenem based antimicrobial compounds, and include for example hydrogen atom or the protective or substituent groups for the amino group to be described in the following:

Namely, preferred examples of the protective and substituent groups for the amino group include alkyl,

alkenyl, aralkyl, aryl, acyl, amide and silyl groups and halogen atoms, wherein as the alkyl groups, there may be mentioned linear or branched lower alkyl groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, tert-butyl and hexyl groups and monocyclic or polycyclic alkyl groups such as cyclopentyl, cyclohexyl, menthyl, fenchyl and bornyl groups; examples of the alkenyl groups include linear or branched lower alkenyl groups such as vinyl, allyl, 1-propenyl, 2-butenyl and 2-methyl-2-propenyl groups; as the aralkyl groups, there may be mentioned, for example, aralkyl groups having a number of carbon atoms of 7 to 24 such as benzyl, phenethyl, trityl and benzhydryl groups; and examples of the aryl groups include aryl groups having a number of carbon atoms of 6 to 10 such as phenyl, tolyl, xylyl, mesityl and cumenyl groups.

As the silyl groups, there may be mentioned tri-substituted silyl groups, which specifically include trialkylsilyl, aryl(alkyl)alkoxysilyl, alkoxydiarylsilyl, triarylsilyl, alkyl-diarylsilyl, aryldialkylsilyl and triaralkylsilyl groups, such as trimethylsilyl, triethylsilyl, triisopropylsilyl, dimethylhexylsilyl, tert-butyl-dimethylsilyl, methyl-diisopropylsilyl, isopropyl-dimethylsilyl, tert-butylmethoxyphenylsilyl, tert-butoxy-diphenylsilyl, triphenylsilyl, tert-butyl-diphenylsilyl, dimethylcumylsilyl and tribenzylsilyl groups.

These alkyl, alkenyl, aralkyl, aryl and silyl groups may be substituted in their individual groups with one or more of substituents, for example, halogen atoms such as fluorine, chlorine and bromine atoms; carboxyl groups; formyl groups; nitro group; cyano group; hydroxyl group; amino groups; linear or branched lower alkyl groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, tert-butyl and hexyl groups; monocyclic or polycyclic alkyl groups such as cyclopentyl, cyclohexyl, menthyl, fenchyl and bornyl groups; linear or branched lower alkenyl groups such as vinyl, allyl, 2-chloroallyl, 1-propenyl, 2-butenyl and 2-methyl-2-propenyl groups; aryl groups having a number of carbon atoms of 6 to 10 such as phenyl, tolyl, xylyl, mesityl and cumenyl groups; aralkyl groups having a number of carbon atoms of 7 to 24

5
10
15

20
25
30
35

aralkyl groups (as is the same with the aralkylthio and
aralkyloxy groups) include for example halogen atoms, and
carboxyl, formyl, nitro, cyano, hydroxyl, amino, alkyl,
alkenyl, aryl, aralkyl, alkylthio, alkenylthio, aralkylthio,
5 arylthio, alkyloxy, alkenyloxy, aralkyloxy, aryloxy,
alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl,
aralkylsulfonyl, arylsulfinyl, arylsulfonyl, carbamoyl,
carbamoyloxy, imino-lower-alkyl, imino-lower-alkylamino,
acyloxy, silyloxy, heterocyclic, heterocyclic-thio,
10 heterocyclic-oxy, acyl, ester, thioester and amide groups;
as the substituents for the aryl groups (as is the same with
the arylthio and aryloxy groups), there may be mentioned
halogen atoms, and carboxyl, formyl, nitro, cyano, hydroxyl,
amino, alkyl, alkenyl, aryl, aralkyl, alkylthio, alkenylthio,
15 aralkylthio, arylthio, alkyloxy, alkenyloxy, aralkyloxy,
aryloxy, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl,
aralkylsulfonyl, arylsulfinyl, arylsulfonyl, carbamoyl,
carbamoyloxy, imino-lower-alkyl, imino-lower-alkylamino,
acyloxy, silyloxy, heterocyclic, heterocyclic-thio,
20 heterocyclic-oxy, acyl, ester, thioester and amide groups;
and the substituents for the amino, carbamoyl, carbamoyloxy,
imino-lower-alkyl, imino-lower-alkylamino and amide groups
include for example halogen atoms, and carboxyl, formyl,
nitro, cyano, hydroxyl, amino, alkyl, alkenyl, aryl, aralkyl,
25 alkylthio, alkenylthio, aralkylthio, arylthio, alkyloxy,
alkenyloxy, aralkyloxy, aryloxy, alkylsulfinyl,
alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl,
arylsulfinyl, arylsulfonyl, carbamoyl, carbamoyloxy, imino-
lower-alkyl, imino-lower-alkylamino, acyloxy, silyloxy,
30 heterocyclic, heterocyclic-thio, heterocyclic-oxy, acyl,
ester, thioester and amide groups.

As the acyl group, there may be mentioned alkyl-
carbonyl, alkenylcarbonyl, aralkylcarbonyl, arylcarbonyl,
alkylthiocarbonyl, alkenylthiocarbonyl, aralkylthiocarbonyl,
35 arylthiocarbonyl, alkyloxycarbonyl, alkenyloxycarbonyl,
aralkyloxycarbonyl, aryloxycarbonyl, silyloxycarbonyl,
heterocyclic-carbonyl, heterocyclic-thiocarbonyl,
heterocyclic-oxycarbonyl, estercarbonyl, thioestercarbonyl
and amidocarbonyl groups which correspond individually to the

above-mentioned alkyl, alkenyl, aralkyl, aryl, alkylthio, alkenylthio, aralkylthio, arylthio, alkyloxy, alkenyloxy, aralkyloxy, aryloxy, silyloxy, heterocyclic, heterocyclic-thio, heterocyclic-oxy, ester and thioester groups as well as
5 the below-described amide groups.

As the amide groups, furthermore, there may be mentioned amide groups which individually may be substituted with one or more of substituents such as the above-mentioned substituents for the amide groups, and examples of the
10 halogen atoms include fluoroine, chlorine and bromine atoms.

The protective groups for the amino group are not particularly limited, and there may be suitably selected and used the conventionally employed protective groups, preferred examples of which are exemplified by the above-mentioned
15 silyl groups; lower-alkoxy lower alkyl groups which may have at least one suitable substituent, such as methoxymethyl and methoxyethoxymethyl groups; aralkyl-oxycarbonyl-lower-alkyl groups which may have at least one suitable substituent, such as benzyloxycarbonylmethyl, p-nitrobenzyloxycarbonylmethyl,
20 p-methoxybenzyloxycarbonylmethyl, phenethyloxycarbonylmethyl, trityloxycarbonylmethyl, benzhydryloxycarbonylmethyl, bis(methoxyphenyl)methyloxycarbonylmethyl, 3,4-dimethoxybenzyloxycarbonylmethyl and 4-hydroxy-3,5-di-tert-butylbenzyloxycarbonylmethyl groups; aralkyl groups which may
25 have at least one suitable substituent, such as benzyl, p-methoxybenzyl, p-nitro-benzyl, p-tert-butylbenzyl, 3,4-dimethylbenzyl, phenethyl, benzhydryl, trityl, bis(methoxyphenyl)methyl, 2,4-dimethoxybenzyl, 3,4-dimethoxybenzyl, 4-hydroxy-3,5-di-tert-butylbenzyl and 2-naphthylmethyl groups; aralkylcarbonyl groups which may have
30 at least one suitable substituent, such as benzylcarbonyl, p-methoxybenzylcarbonyl, p-nitrobenzylcarbonyl, p-tert-butylbenzylcarbonyl, 3,4-dimethylbenzylcarbonyl, phenethylcarbonyl, benzhydrylcarbonyl, tritylcarbonyl,
35 bis(methoxyphenyl)methylcarbonyl, 2,4-dimethoxybenzylcarbonyl, 3,4-dimethoxybenzylcarbonyl, 4-hydroxy-3,5-di-tert-butylbenzylcarbonyl and 2-naphthylmethylcarbonyl groups; heterocyclic groups which may have at least one suitable substituent, such as tetrahydropyranyl group; arylcarbonyl

groups which may have at least one suitable substituent, such as benzoyl, chlorobenzoyl, p-methoxybenzoyl, p-nitrobenzoyl, p-tert-butylbenzoyl, toluoyl and naphthoyl groups; arylcarbonyl-lower-alkyl groups which may have at least one
5 suitable substituent, such as benzoylmethyl, chlorobenzoylmethyl, p-methoxybenzoylmethyl, p-nitrobenzoylmethyl, p-tert-butylbenzoylmethyl, toluoylmethyl and naphthoylmethyl groups; aryloxy-lower-alkylcarbonyl groups which may have at least one suitable substituent, such as phenoxyacetyl, 4-
10 chlorophenoxyacetyl, tolyloxyacetyl, tert-butylphenoxyacetyl, xylyloxyacetyl, mesityloxyacetyl and cumenyloxyacetyl groups; arylsulfonyl groups which may have at least one suitable substituent, such as benzenesulfonyl, p-tert-butylbenzenesulfonyl and toluenesulfonyl groups; alkyl-
15 sulfonyl groups which may have at least one suitable substituent, such as mesyl group; formyl group; aliphatic carboxylic acid acyl groups which may have at least one suitable substituent, such as acetyl, chloroacetyl, bromoacetyl, dichloroacetyl, trichloroacetyl, methoxyacetyl, propionyl, butyryl, isobutyryl, valeryl, pivaloyl, hexanoyl,
20 2-ethylbutyryl, 3,3-dimethyl-butyl, pentanoyl, caprylyl, decanoyl and acryloyl groups; lower-alkoxycarbonyl groups which may have at least one suitable substituent, such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, tert-butoxycarbonyl, 2-iodoethoxy-
25 carbonyl, 2,2-dichloroethoxycarbonyl and 2,2,2-trichloroethoxycarbonyl groups; lower-alkenyloxycarbonyl groups which may have at least one suitable substituent, such as vinylloxycarbonyl, allyloxycarbonyl and 2-chloroallyl-
30 oxycarbonyl groups; aralkyloxycarbonyl groups which may have at least one suitable substituent, such as benzyloxycarbonyl, p-nitrobenzyloxycarbonyl, benzhydryloxycarbonyl and trityloxycarbonyl groups; aryloxycarbonyl groups which may have at least one suitable substituent, such as
35 phenoxycarbonyl, 4-chlorophenoxycarbonyl and tert-butylphenoxycarbonyl groups; and carbamoyl groups which may have at least one suitable substituent and their corresponding thiocarbamoyl groups, such as methylcarbamoyl, phenylcarbamoyl and naphthylcarbamoyl groups.

Examples

~~Examples~~

Below described are examples to illustrate the present invention in more detail, but this invention is not understood to be limited to these examples.

Example 1

Synthesis of (3S,4S)-3-[(R)-1-(tert-butyldimethylsilyloxy)-ethyl]-4-di(methoxycarbonyl)methyl-2-azetidinone

A solution of dimethyl malonate (291 mg, 2.2 mmole) in tetrahydrofuran (5 ml) was added to a suspension of sodium hydride (90 mg, 2.2 mmole) in tetrahydrofuran (10 ml) in a nitrogen stream, while stirring under ice cooling, and the solution mixture was stirred for 10 min. and then admixed with cuprous bromide dimethylsulfide complex (452 mg, 2.2 mmole), followed by stirring for 15 min, addition of a solution of (3S,4R)-3-[(R)-1-(tert-butyldimethylsilyloxy)-ethyl]-4-phenylthio-2-azetidinone (338 mg, 1 mmole) in tetrahydrofuran (10 ml) and stirring for 15 min. The reaction solution was admixed with saturated ammonium chloride solution, and after the insoluble matters were removed by filtration, the filtrate was extracted with ethyl acetate. The ethyl acetate layer was washed with aqueous saturated sodium chloride solution three times, dried over magnesium sulfate and concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (the developing solvent was a 7:1 mixture of n-hexane and ethyl acetate) to give 267 mg (74.4 %) of the subject compound in the form of colorless solid.

IR (KBr) cm^{-1} : 3158, 3103, 2958, 1770, 1774, 1440, 1202, 836. NMR (CDCl_3): δ 0.07(6H,s,-Si(CH₃)₂), 0.88(9H,s,-SiC(CH₃)₃), 1.11(3H,d,J=6.6Hz, CH₃CH-OSi), 3.04(1H,m,C3-H), 3.55(1H,d,J=9.2Hz, CH(CO₂CH₃)₂), 3.77 and 3.78(each 3H,s, CO₂CH₃), 4.18-4.28(1H,m,CH₃CH-OSi), 6.06(1H,brs,-NH)

Example 2

Synthesis of (3S,4S)-3-[(R)-1-(tert-butyldimethylsilyloxy)-ethyl]-4-[1,1-di(ethoxycarbonyl)ethyl]-2-azetidinone

A solution of diethyl 2-methylmalonate (192 mg, 1.1 mmole) in tetrahydrofuran (2.5 ml) was added to a suspension of sodium hydride (44 mg, 1.1 mmole) in tetrahydrofuran (5 ml) in a nitrogen stream, while stirring under ice cooling,

and the solution mixture was stirred for 10 min. and then admixed with cuprous bromide dimethylsulfide complex (227 mg, 1.1 mmole), followed by stirring for 15 min, addition of a solution of (3S,4R)-3-[(R)-1-(tert-butyldimethylsilyloxy)ethyl]-4-phenylthio-2-azetidinone (169 mg, 0.5 mmole) in tetrahydrofuran (5 ml) and stirring for 15 min. The reaction solution was admixed with saturated ammonium chloride solution, and after the insoluble matters were removed by filtration, the filtrate was extracted with ethyl acetate. The ethyl acetate layer was washed with aqueous saturated sodium chloride solution three times, dried over magnesium sulfate and concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (the developing solvent was a 7:1 mixture of n-hexane and ethyl acetate) to give 106 mg (52.8 %) of the subject compound in the form of colorless solid.

IR (KBr) cm^{-1} : 3178, 2856, 1769, 1756, 1736, 1256, 1114, 837. NMR (CDCl_3): δ 0.07(6H,s,-Si(CH₃)₂), 0.88(9H,s,-SiC(CH₃)₃), 1.14(3H,d,J=5.9Hz, CH₃CH-OSi), 1.23-1.31(6H,m,(CO₂CH₂CH₃)₂), 1.46(3H,s,CCH₃(CO₂CH₂CH₃)₂), 3.01(1H,d,J=1.92Hz,C4-H), 4.12-4.28(6H,m,CH₃CH-OSi, C3-H and (CO₂CH₂CH₃)₂), 5.97(1H,brs,-NH)

Example 3

Synthesis of (3S,4S)-3-[(R)-1-(tert-butyldimethylsilyloxy)-ethyl]-4-di(ethoxycarbonyl)fluoromethyl-2-azetidinone

A solution of diethyl 2-fluoromalonate (196 mg, 1.1 mmole) in tetrahydrofuran (2.5 ml) was added to a suspension of sodium hydride (44 mg, 1.1 mmole) in tetrahydrofuran (5 ml) in a nitrogen stream, while stirring under ice cooling, and the solution mixture was stirred for 10 min. and then admixed with cuprous bromide dimethylsulfide complex (227 mg, 1.1 mmole), followed by stirring for 15 min, addition of a solution of (3S,4R)-3-[(R)-1-(tert-butyldimethylsilyloxy)ethyl]-4-phenylthio-2-azetidinone (169 mg, 0.5 mmole) in tetrahydrofuran (5 ml) and stirring for 15 min. The reaction solution was admixed with saturated ammonium chloride solution, and after the insoluble matters were removed by filtration, the filtrate was extracted with ethyl acetate. The ethyl acetate layer was washed with aqueous

saturated sodium chloride solution three times, dried over magnesium sulfate and concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (the developing solvent was a 7:1 mixture of n-hexane and ethyl acetate) to give 166 mg (82 %) of the subject compound in the form of colorless solid.

IR (KBr) cm^{-1} : 3169, 3104, 2854, 1744, 1473, 1289, 1251, 836. NMR (CDCl_3): δ 0.07(6H,s,-Si(CH₃)₂), 0.88(9H,s,-SiC(CH₃)₃), 1.08(3H,d,J=5.9Hz, CH₃CH-OSi), 1.30-1.36 (6H,m,(CO₂CH₂CH₃)₂), 3.30(1H,s,C4-H), 4.24-4.45(6H,m,CH₃CH-OSi, C3-H and (CO₂CH₂CH₃)₂), 5.94(1H,brs,-NH)

Example 4

Synthesis of (3S,4S)-3-[(R)-1-(tert-butyldimethylsilyloxy)-ethyl]-4-(hydroxycarbonyl)fluoromethyl-2-azetidinone

Dissolved in 1 ml of pyridine was (3S,4S)-3-[(R)-1-(tert-butyldimethylsilyloxy)ethyl]-4-di(ethoxycarbonyl)-fluoromethyl-2-azetidinone (405 mg, 1.0 mmole), and 0.5 ml of 4N sodium hydroxide solution was added to the solution, followed by stirring at room temperature for 1.5 hours. The reaction solution was admixed with 0.4 ml of 5N hydrochloric acid and 2.0 ml of 2,4,6-collidine, followed by heating at 150°C and stirring for 1 hour. After the reaction solution was allowed to cool, the solvent was distilled off, and the residue was dissolved in chloroform. The solution was washed with aqueous saturated potassium hydrogensulfate solution twice, admixed with aqueous saturated sodium hydrogen-carbonate solution and separated into two layers. And the aqueous layer was washed with chloroform once, admixed with potassium hydrogensulfate and sodium chloride, and extracted with ethyl acetate. The ethyl acetate layer was washed with aqueous saturated sodium chloride solution twice, dried over magnesium sulfate and concentrated under reduced pressure, and the residue was recrystallized from ethyl acetate/hexane to give 254 mg (83 %) of the subject compound consisting of a diastereomer mixture at a ratio of about 2:1 in the form of colorless solid.

IR (KBr) cm^{-1} : 3406, 2929, 1748, 1473, 1376, 1254, 1168, 1143, 1112, 1072, 1041, 962, 839, 780. NMR ($\text{DMSO}-d_6$) major product: δ 0.02 and 0.05(each 3H,s,-Si(CH₃)₂), 0.84(9H,s,-

SiC(CH₃)₃), 1.05(3H,d,J=5.9Hz, CH₃CH-OSi), 3.08(1H,m,C3-H), 3.88(1H,m,J=21Hz,C4-H), 4.13(1H,m,CH₃CH-OSi), 5.20(1H,dd,J=49Hz,2.6Hz,F-CH-COOH), 8.11(1H,brs,-NH), 13.62(1H,brs,-COOH). By-product: δ 0.03 and 0.05(each 3H,s,-Si(CH₃)₂), 0.84(9H,s,-SiC(CH₃)₃), 1.11(3H,d,J=6.6Hz, CH₃CH-OSi), 3.08(1H,m,C3-H), 3.88(1H,m,J=21Hz,C4-H), 4.13(1H,m,CH₃CH-OSi), 5.08(1H,dd,J=48Hz,4.6Hz,F-CH-COOH), 8.29(1H,brs,-NH), 13.62(1H,brs,-COOH). MS(FAB) m/z = 306(C₁₃H₂₄NO₄FSi+H)⁺

10 Example 5

Synthesis of (3S,4S)-3-[(R)-1-(tert-butyldimethylsilyloxy)-ethyl]-4-(1-acetyl-1-ethoxycarbonyl-ethyl)-2-azetidinone

Ethyl 2-methylacetoacetate (0.87 ml, 6.03 mmole) was added to a suspension of sodium hydride (240 mg, 6.0 mmole) in tetrahydrofuran (5 ml) in a nitrogen stream, and the solution mixture was stirred for 10 min at room temperature and admixed with cuprous bromide dimethylsulfide complex (1.23 g, 5.98 mmole), followed by stirring for 1 hour, addition of (3S,4R)-3-[(R)-1-(tert-butyldimethylsilyloxy)ethyl]-4-phenylthio-2-azetidinone (1.01 g, 2.99 mmole) and stirring for 3 hours. The reaction solution was admixed with saturated ammonium chloride solution, and after the insoluble matters were removed by filtration, the filtrate was extracted with ethyl acetate. The ethyl acetate layer was washed with aqueous saturated sodium chloride solution three times, dried over magnesium sulfate and concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (the developing solvent was a 4:1 mixture of n-hexane and ethyl acetate) to give 876 mg (78.9 %) of the subject compound consisting of a diastereomer mixture at a ratio of about 5:4 in the form of colorless solid.

IR (KBr) cm⁻¹: 3178, 2957, 2929, 2856, 1767, 1744, 1716, 1646, 1472, 1377, 1362, 1340, 1250, 1159, 1105, 1063, 967, 837, 777. NMR (CDCl₃): Major product: δ 0.06(6H,s,-Si(CH₃)₂), 0.88(9H,s,-SiC(CH₃)₃), 1.19(3H,d,J=5.9Hz, CH₃CH-OSi), 1.27(3H,t,J=7.2Hz,CO₂CH₂CH₃), 1.42(3H,s,CCH₃(CO₂CH₂CH₃)-(COCH₃)), 2.21(3H,s,CCH₃(CO₂CH₂CH₃)(COCH₃)), 2.91

(1H,m,C3-H), 4.03(1H,d,J=2.0Hz,C4-H), 4.16-4.26(3H,m,CH₃CH-OSi and CO₂CH₂CH₃), 5.89(1H,brs,-NH). By-product: δ 0.07 (6H,s,-Si(CH₃)₂), 0.88(9H,s,-SiC(CH₃)₃), 1.06(3H,d,J=6.6Hz, CH₃CH-OSi), 1.31(3H,t,J=7.2Hz,CO₂CH₂CH₃), 1.39(3H,s,CCH₃-(CO₂CH₂CH₃)(COCH₃)), 2.18(3H,s,CCH₃(CO₂CH₂CH₃)(COCH₃)), 2.93(1H,m,C3-H), 4.16-4.26(4H,m,CH₃CH-OSi, C4-H and CO₂CH₂CH₃), 5.89(1H,brs,-NH)

Example 6

Synthesis of (3S,4S)-3-[(R)-1-(tert-butyldimethylsilyloxy)-ethyl]-4-(1-acetyl-1-allyloxycarbonyl-ethyl)-2-azetidinone

Allyl 2-methylacetoacetate (624 mg, 4.00 mmole) was added to a suspension of sodium hydride (160 mg, 4.0 mmole) in tetrahydrofuran (10 ml) in a nitrogen stream, and the solution mixture was stirred for 10 min at room temperature and then admixed with cuprous bromide dimethylsulfide complex (822 mg, 5.98 mmole), followed by stirring for 1 hour, addition of (3S,4R)-3-[(R)-1-(tert-butyldimethylsilyloxy)ethyl]-4-phenylthio-2-azetidinone (677 mg, 2.00 mmole) and stirring for 3 hours. The reaction solution was admixed with saturated ammonium chloride solution, and after the insoluble matters were removed by filtration, the filtrate was extracted with ethyl acetate. The ethyl acetate layer was washed with aqueous saturated sodium chloride solution three times, dried over magnesium sulfate and concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (the developing solvent was a 3:1 mixture of n-hexane and ethyl acetate) to give 495 mg (64.5 %) of the subject compound consisting of a diastereomer mixture (Products A and B) at a ratio of about 1:1 in the form of colorless solid.

IR (KBr) cm⁻¹: 3178, 2956, 2929, 2857, 1769, 1715, 1250, 1158, 1104, 1063, 966, 838, 778. NMR (CDCl₃): Product A: δ 0.06(6H,s,-Si(CH₃)₂), 0.88(9H,s,-SiC(CH₃)₃), 1.19(3H,d,J=6.6Hz, CH₃CH-OSi), 1.44(3H,s,CCH₃(CO₂)(COCH₃)), 2.21(3H,s,CCH₃(CO₂)(COCH₃)), 2.92(1H,m,C3-H), 4.15-4.24(1H,m,CH₃CH-OSi), 4.25(1H,d,J=2.0Hz,C4-H), 4.62-4.69(2H,m,OCH₂CHCH₂), 5.28-5.40(2H,m,OCH₂CHCH₂), 5.83-5.93(2H,m,-NH and OCH₂CHCH₂). Product B: δ 0.06 (6H,s,-Si(CH₃)₂), 0.88(9H,s,-SiC(CH₃)₃), 1.06(3H,d,J=5.9Hz, CH₃CH-

OSi), 1.40(3H,s,CCH₃(CO₂)(COCH₃)), 2.18(3H,s,
CCH₃(CO₂)(COCH₃)), 2.94(1H,m,C3-H), 4.05(1H,d,J=2.6Hz, C4-H),
4.15-4.24(1H,m,CH₃CH-OSi) 4.62-4.69(2H,m,OCH₂CHCH₂), 5.28-
5.40(2H,m,OCH₂CHCH₂), 5.83-5.93(2H,m,-NH and OCH₂CHCH₂).

5 Example 7

Synthesis of (3S,4S)-3-[(R)-1-(tert-butyldimethylsilyloxy)-
ethyl]-4-(1-acetyl-1-benzoyloxycarbonyl-ethyl)-2-azetidinone

Benzyl 2-methylacetoacetate (825 mg, 4.00 mmole) was
added to a suspension of sodium hydride (160 mg, 4.0 mmole)
10 in tetrahydrofuran (10 ml) in a nitrogen stream, and the
solution mixture was stirred for 10 min at room temperature
and then admixed with cuprous bromide dimethylsulfide
complex (822 mg, 5.98 mmole), followed by stirring for 1
hour, addition of (3S,4R)-3-[(R)-1-(tert-butyldimethyl-
15 silyloxy)ethyl]-4-phenylthio-2-azetidinone (677 mg, 2.00
mmole) and stirring for 3 hours. The reaction solution was
admixed with saturated ammonium chloride solution, and after
the insoluble matters were removed by filtration, the
filtrate was extracted with ethyl acetate. The ethyl acetate
20 layer was washed with aqueous saturated sodium chloride
solution three times, dried over magnesium sulfate and
concentrated under reduced pressure, and the residue was
purified by silica gel column chromatography (the developing
solvent was a 3:1 mixture of n-hexane and ethyl acetate) to
25 give 673 mg (77.6 %) of the subject compound consisting of a
diastereomer mixture (Products A and B) at a ratio of about
1:1 in the form of colorless oily matter.

IR (neat) cm⁻¹: 3259, 2955, 2930, 2856, 1770, 1713,
1498, 1462, 1455, 1373, 1360, 1253, 1142, 1104, 1067, 960,
30 837, 778, 695. NMR (CDCl₃): Product A: δ 0.05(6H,s,-
Si(CH₃)₂), 0.87(9H,s,-SiC(CH₃)₃), 1.03 (3H,d,J=6.6Hz, CH₃CH-
OSi), 1.39(3H,s,CCH₃(CO₂)(COCH₃)), 2.04(3H,s,CCH₃(CO₂)-
(COCH₃)), 2.92(1H,m,C3-H), 4.11-4.22(1H,m,CH₃CH-OSi),
4.27(1H,d,J=2.0Hz,C4-H), 5.10-5.28(2H,m,OCH₂-C₆H₅),
35 5.83(1H,brs,-NH), 7.30-7.38(5H,m,C₆H₅). Product B: δ 0.05 and
0.07(each 3H,s,-Si(CH₃)₂), 0.88(9H,s,-SiC(CH₃)₃),
1.17(3H,d,J=6.6Hz, CH₃CH-OSi), 1.43(3H,s,CCH₃(CO₂)(COCH₃)),
2.12(3H,s,CCH₃(CO₂)(COCH₃)), 2.93(1H,m,C3-H), 4.06(1H,d,

J=2.0Hz, C4-H), 4.11-4.22(1H, m, CH₃CH-OSi), 5.10-5.28 (2H, m, OCH₂C₆H₅), 5.83(1H, brs, -NH), 7.30-7.38(5H, m, C₆H₅)

Example 8

Synthesis of (3S,4S)-3-[(R)-1-(tert-butyldimethylsilyloxy)-ethyl]-4-(1-acetyloethyl)-2-azetidinone

(3S,4S)-3-[(R)-1-(tert-Butyldimethylsilyloxy)ethyl]-4-(1-acetyl-1-allyloxycarbonyl-ethyl)-2-azetidinone (192 mg, 0.50 mmol) was dissolved in tetrahydrofuran (10 ml) under a nitrogen stream, and dimedone (70 mg, 0.50 mmol) and tetrakis(triphenylphosphine)palladium (58 mg, 0.05 mmol) were added to the resultant solution, followed by stirring under reflux heating for 3 hours. The reaction solution was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (the developing solvent was a 1:1 mixture of n-hexane and ethyl acetate) to give 108 mg (72.0 %) of the subject compound consisting of a diastereomer mixture at a ratio of about 5:1 in the form of colorless solid.

IR (KBr) cm⁻¹: 3314, 2954, 1717, 1713, 1654, 1617, 1437, 1308, 1276, 1170, 840. NMR (CDCl₃): Major product: δ 0.07 and 0.08 (each 3H, s, -Si(CH₃)₂), 0.88(9H, s, -SiC(CH₃)₃), 1.23 (3H, d, J=7.3Hz, CHCH₃(COCH₃)), 1.25(3H, d, J=6.0Hz, CH₃CH-OSi), 2.20(3H, s, CHCH₃(COCH₃)), 2.58(1H, dq, J=9.9Hz and 7.6Hz, CHCH₃(COCH₃)), 2.72(1H, dd, J=1.3Hz and 5.9Hz, C3-H), 3.67(1H, dd, J=2.0Hz and 9.9Hz, C4-H), 4.16(1H, m, CH₃CH-OSi), 5.98(1H, brs, -NH).

By-product B: δ 0.07(6H, s, -Si(CH₃)₂), 0.88(9H, s, -SiC(CH₃)₃), 1.18 (3H, d, J=7.2Hz, CHCH₃(COCH₃)), 1.20(3H, d, J=6.6Hz, CH₃CH-OSi), 2.22(3H, s, CHCH₃(COCH₃)), 2.81(1H, m, CHCH₃(COCH₃)), 2.87(1H, dd, J=2.0Hz and 5.3Hz, C3-H), 3.89(1H, dd, J=2.0Hz and 4.6Hz, C4-H), 4.16(1H, m, CH₃CH-OSi), 5.92(1H, brs, -NH), MS (FAB) m/z = 300(C₁₅H₂₉NO₃Si+H)⁺

Example 9

Synthesis of (3S,4S)-3-[(R)-1-(tert-butyldimethylsilyloxy)-ethyl]-4-ethoxycarbonylmethyl-2-azetidinone

A mixture of (3S,4R)-3-[(R)-1-(tert-butyldimethylsilyloxy)ethyl]-4-phenylthio-2-azetidinone (338 mg, 1 mmol), zinc (260 mg, 4 mg atom), cuprous bromide dimethylsulfide complex (411 mg, 2 mmol), ethyl bromoacetate (0.23 ml, 2

mmol) and tetrahydrofuran (4.5 ml) was stirred at room temperature for 31 hours. The reaction mixture was admixed with saturated ammonium chloride solution and extracted with ethyl acetate, and the ethyl acetate layer was washed with aqueous saturated sodium chloride solution three times, dried over magnesium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (the developing solvent was a 7:1 mixture of n-hexane and ethyl acetate) to give 90 mg (29 %) of the subject compound in the form of colorless solid.

IR (KBr) cm^{-1} : 3186, 2954, 1761, 1729, 1472, 1248, 1140.
NMR (CDCl_3): δ 0.08(6H,s,-Si(CH₃)₂), 0.88(9H,s,-SiC(CH₃)₃), 1.21(3H,d,J=6.6Hz, CH₃CH-OSi), 2.56(1H,dd,J=9,9Hz and 16.5Hz,-CH₂CO₂-), 2.73(1H,dd,J=4.0Hz and 16.5Hz,-CH₂CO₂-), 2.82(1H,dd,J=2.0Hz and 4.6Hz,C3-H), 3.97(1H,m,C4-H), 4.13-4.24(3H,m,CH₃CH-OSi), 6.15(1H,brs,-NH)

Example 10

(3S,4S)-3-[(R)-1-(tert-Butyldimethylsilyloxy)ethyl]-4-[1,1-di(allyloxy)carbonyl]ethyl]-2-azetidinone

A solution of diallyl 2-methylmalonate (396 mg, 2 mmol) in tetrahydrofuran (2 ml) was added dropwise to a suspension of sodium hydride (80 mg, 2 mmol) in tetrahydrofuran (5 ml) under an argon stream at 0°C, and the solution mixture was stirred for 5 min and admixed with cuprous bromide dimethylsulfide complex (411 mg, 2 mmol), followed by stirring for 30 min at the same temperature, addition of a solution of (3R,4S)-3-[(R)-1-(tert-butyldimethylsilyloxy)ethyl]-4-phenylthio-2-azetidinone (338 mg, 1 mmole) in tetrahydrofuran (3 ml) and stirring for 1 hour. The reaction solution was admixed with aqueous saturated ammonium chloride solution and extracted with ethyl acetate. The ethyl acetate layer was dried over magnesium sulfate and concentrated under reduced pressure, and the residue was purified by column chromatography (10 g of silica gel; a 8/1 mixture of n-hexane/ethyl acetate) to give 392 mg (92 %) of the subject compound in the form of colorless crystals.

NMR (CDCl_3): δ 0.07(6H,s,-Si(CH₃)₂), 0.88(9H,s,SiC(CH₃)₃), 1.13(3H,d,J=6.6Hz, CH₃CHOSi), 1.50 (3H,s,CH₃), 3.01-3.04(1H,m,C3-H), 4.18(1H,d,J=2.6Hz,C4-H), 4.17-4.25

(1H,m,CHOSi), 4.60-4.68(4H,m,CH₂CH=CH₂), 5.23-5.38(4H,m,
CH₂CH=CH₂), 5.77-5.95(2H,m,CH₂CH=CH₂), 5.95(1H,brs,-NH)

Example 11

5 (3S,4S)-3-[(R)-1-(tert-Butyldimethylsilyloxy)ethyl]-4-[(RS)-
1-allyloxycarbonyl-cyclohexan-2-on-1-yl]-2-azetidinone

10 A solution of 2-allyloxycarbonylcyclohexanone (1.27 g,
6.98 mmol) in tetrahydrofuran (8 ml) was added dropwise to a
suspension of sodium hydride (267 mg, 6.98 mmol) in
tetrahydrofuran (30 ml) under an argon stream at 0°C, and the
15 solution mixture was stirred for 10 min and admixed with
cuprous bromide dimethylsulfide complex (1.44 g, 6.98 mmol),
followed by stirring for 15 min at the same temperature,
addition of a solution of (3R,4S)-3-[(R)-1-(tert-
butyldimethylsilyloxy)ethyl]-4-phenylthio-2-azetidinone (1.18
20 g, 3.49 mmol) in tetrahydrofuran (8 ml) and stirring for 40
min. The reaction solution was admixed with aqueous saturated
ammonium chloride solution and extracted with ethyl acetate.
The ethyl acetate layer was washed with aqueous saturated
sodium chloride solution, dried over magnesium sulfate and
25 concentrated under reduced pressure, and the residue was
purified by column chromatography (50 g of silica gel; a 4/1
mixture of n-hexane/ethyl acetate) to give 340 mg (24 %) of
the subject compound having the cyclohexanone ring in the R-
configuration in the form of colorless crystals and 920 mg
30 (64 %) of the subject compound having the cyclohexanone ring
in the S-configuration in the form of colorless oily
substance.

R-configuration isomer: IR (KBr) cm⁻¹: 1718, 1767. NMR
(CDCl₃): δ 0.06 and 0.07(total 6H,each s,Si(CH₃)₂), 0.87
35 (9H,s,SiC(CH₃)₃), 1.00(3H,d,J=5.9Hz,CH₃CHOSi), 1.51-1.90
(4H,m), 2.01-2.12(1H,m), 2.36-2.54(3H,m), 2.95-3.00(1H,m,C3-
H), 4.17-4.28(1H,m,CHOSi), 4.37(1H,d,J=2.0Hz, C4-H), 4.57-
4.70(2H,m,CH₂CH=CH₂), 5.26-5.42(2H,m, CH₂CH=CH₂),
5.67(1H,brs,-NH), 5.82-5.98(1H,m,CH₂CH=CH₂)
S-configuration isomer: IR (neat) cm⁻¹: 1711, 1766. NMR
(CDCl₃): δ 0.06 and 0.07(total 6H,each s,Si(CH₃)₂), 0.87
(9H,s,SiC(CH₃)₃), 1.20(3H,d,J=6.6Hz,CH₃CHOSi), 1.39-1.70
(3H,m), 1.75-1.88(1H,m), 2.00-2.11(1H,m), 2.42-2.52(3H,m),

3.12-3.17 (1H,m,C3-H), 3.88(1H,d,J=2.0Hz,C4-H), 4.10-4.22 (1H,m, CHOSi), 4.63-4.68(2H,m,CH₂CH=CH₂), 5.25-5.37(2H,m, CH₂CH=CH₂), 5.78-5.92(1H,m,CH₂CH=CH₂), 5.93(1H,brs,-NH)

Example 12

5 (3S,4S)-3-[(R)-1-(tert-Butyldimethylsilyloxy)ethyl]-4-[(RS)-1-(hydroxycarbonyl)ethyl]-2-azetidinone

Triethylamine (60 µl, 0.430 mmol), formic acid (15 µl, 0.401 mmol), triphenylphosphine (3.8 mg, 0.014 mmol) and palladium acetate (1.6 mg, 0.007 mmol) were added to a
10 solution of the compound (61 mg, 0.143 mmol) as obtained in Example 10 in 1,4-dioxane (1.5 ml), successively, under an argon stream, and the reaction mixture was stirred at 100°C for one and a half hours. The mixture was cooled and admixed with diethyl ether (7 ml) and 0.5N sodium hydroxide solution
15 (7 ml), followed by stirring, and the aqueous layer was adjusted to pH 3 with 1N hydrochloric acid and extracted with ether (30 ml). The extract layer was dried over sodium sulfate and concentrated under reduced pressure to give 36 mg (83 %) of the subject compound as a mixture consisting of the
20 α(R-isomer)/β(S-isomer) at a ratio of 3/1.

NMR (CDCl₃): δ 0.07 and 0.09(total 6H,s,Si(CH₃)₂), 0.88(9H,s,SiC(CH₃)₃), 1.15-1.31(6H,m,CH₃CHOSi and CH₂CHCO₂H), 2.55(0.75H,dd,J=7.3 and 9.9Hz,CHCO₂H), 2.70-2.85(1H,m,CHCO₂H and C3-H), 3.02-3.06(0.25H,m,C3-H), 3.69 (0.75H,dd,J=2.0Hz and 9.9Hz,C4-H), 3.95(0.25H, dd,J=2.0Hz and 4.5Hz,C4-H),
25 4.12-4.25(1H,m,CHOSi), 6.45(0.25H,brs,-NH), 6.79(0.75H,brs, NH)

Example 13

30 (3S,4S)-3-[(R)-1-(tert-Butyldimethylsilyloxy)ethyl]-4-[(RS)-cyclohexan-2-on-1-yl]-2-azetidinone

Under an argon stream, a solution of triethylamine (87 µl, 0.625 mmol) and formic acid (19 µl, 0.50 mmol) in tetrahydrofuran (1 ml) was added dropwise to a solution of triphenylphosphine (6.6 mg, 0.025 mmol) and palladium acetate
35 (2.8 mg, 0.0125 mmol) in tetrahydrofuran (1.5 ml), and then a solution of the S-isomer (102 mg, 0.25 mmol) of the compound as obtained in Example 11 in tetrahydrofuran (0.8 ml) was added dropwise to the resultant solution, followed by stirring at room temperature for 1 hour. The reaction mixture

was admixed with aqueous saturated ammonium chloride solution (5 ml) and extracted with ethyl acetate (30 ml). The ethyl acetate layer was washed with aqueous saturated sodium chloride solution, dried over magnesium sulfate and concentrated under reduced pressure to give the subject compound as a mixture consisting of the α (R-isomer)/ β (S-isomer) at a ratio of 2.5/1. Purification by column chromatography (5 g of silica gel; a 2/1 mixture of hexane/ethyl acetate) yielded 73 mg (90 %) of the subject compound as a mixture.

α -Form; IR (KBr) cm^{-1} : 1710, 1760. NMR (CDCl_3): δ 0.06 and 0.07 (total 6H, each s, $\text{Si}(\text{CH}_3)_2$), 0.88 (9H, s, $\text{SiC}(\text{CH}_3)_3$), 1.22 (3H, d, $J=5.9\text{Hz}$, CH_3CHOSi), 1.32-1.48 (1H, m), 1.61-1.76 (2H, m), 1.86-2.00 (1H, m), 2.06-2.21 (2H, m), 2.23-2.48 (3H, m), 2.70 (1H, dd, $J=2.0$ and 5.9Hz , C3-H), 3.61 (1H, dd, $J=2.0$ and 9.9Hz , C4-H), 4.07-4.22 (1H, m, CHOSi), 6.09 (1H, brs, -NH)

β -Form; IR (KBr) cm^{-1} : 1706, 1755. NMR (CDCl_3): δ 0.06 and 0.07 (total 6H, each s, $\text{Si}(\text{CH}_3)_2$), 0.87 (9H, s, $\text{SiC}(\text{CH}_3)_3$), 1.23 (3H, d, $J=5.9\text{Hz}$, CH_3CHOSi), 1.54-1.81 (3H, m), 1.92-2.20 (3H, m), 2.26-2.48 (2H, m), 2.51-2.61 (1H, m), 2.87 (1H, dd, $J=2.0$ and 4.7Hz , C3-H), 4.06-4.12 (1H, m, C4-H), 4.12-4.24 (1H, m, CHOSi), 5.72 (1H, brs, NH)

Example 14

(3S,4S)-3-[(R)-1-(tert-Butyldimethylsilyloxy)ethyl]-4-[1,1-di(allyloxycarbonyl)ethyl]-1-(p-nitrobenzyloxycarbonyl)-methyl-2-azetidinone

Potassium carbonate (323 mg, 1.61 mmol) was added to a solution of the compound (220 mg, 0.517 mmol) obtained in Example 10 and p-nitrobenzyl 1-iodoacetate (0.592 mmol) in N,N-dimethylformamide (5 ml) under an argon stream, followed by stirring at room temperature for 2 hours. The solution mixture was furthermore stirred at 50°C for 2 hours, cooled, admixed with water (10 ml) and extracted with diethyl ether (45 ml), and the ether layer was washed with aqueous saturated sodium chloride solution (15 ml), dried over magnesium sulfate and concentrated under reduced pressure. The residue was purified by column chromatography (10 g of silica gel, a 20/1 mixture of hexane/ethyl acetate) to give

148 mg (46 %) of the subject compound, with 120 mg (54 %) of the starting compound being recovered.

NMR (CDCl₃): δ 0.02 and 0.06 (total 6H, each s, Si(CH₃)₂), 0.86 (9H, s, SiC(CH₃)₃), 1.20 (3H, d, J=6.0 Hz, CH₃CHOSi), 1.54 (3H, s, CH₃), 3.05 (1H, dd, J=2.0 and 5.9 Hz, C3-H), 3.90 (1H, d, J=17.8 Hz, C(H)HCO₂PNB), 4.08-4.19 (1H, m, CHOSi), 4.29 (1H, d, J=17.8 Hz, C(H)HCO₂PNB), 4.34 (1H, d, J=2.0 Hz, C4-H), 4.53-4.64 (4H, m, CO₂CH₂CH=CH₂), 5.20-5.36 (6H, m, CH₂Ar and CH₂CH=CH₂), 5.75-5.96 (2H, m, CH₂CH=CH₂), 7.55 (2H, d, J=9.2 Hz, Ar), 8.22 (2H, d, J=9.2 Hz, Ar)

Example 15

(3S,4S)-3-[(R)-1-(tert-Butyldimethylsilyloxy)ethyl]-4-[(R)-1-allyloxycarbonyl-cyclohexan-2-on-1-yl]-1-(p-nitrobenzyl-oxycarbonyl)methyl-2-azetidinone

Potassium carbonate (101 mg, 0.733 mmol) was added to a solution of the compound (R-isomer) (100 mg, 0.244 mmol) obtained in Example 11 and p-nitrobenzyl 1-iodoacetate (0.293 mmol) in N,N-dimethylformamide (4 ml) under an argon stream, followed by stirring at room temperature for 12 hours. The solution mixture was furthermore stirred at 50°C for 2 hours, cooled, admixed with water (6 ml) and extracted with diethyl ether (30 ml), and the ether layer was washed with aqueous saturated sodium chloride solution (10 ml), dried over sodium sulfate and concentrated under reduced pressure. The residue was purified by column chromatography (10 g of silica gel, a 20/1 mixture of dichloromethane/ethyl acetate) to give 39 mg (27 %) of the subject compound, with 48 mg (48 %) of the starting compound being recovered.

NMR (CDCl₃): δ 0.06 and 0.07 (total 6H, each s, Si(CH₃)₂), 0.87 (9H, s, SiC(CH₃)₃), 0.99 (3H, d, J=6.5 Hz, CH₃CHOSi), 1.52-1.71 (2H, m), 1.83-1.95 (2H, m), 2.21-2.33 (2H, m), 2.36-2.43 (2H, m), 3.08-3.10 (1H, m, C3-H), 4.09 (1H, d, J=8.7 Hz, C(H)HCO₂PNB), 4.18 (1H, d, J=8.7 Hz, C(H)HCO₂PNB), 4.18-4.25 (1H, m, CHOSi), 4.52 (1H, d, J=2.0 Hz, C4-H), 4.63 (2H, ddd, J=4.5, 8.8 and 13.5 Hz, CO₂CH₂CH=CH₂), 5.13 (1H, d, J=9.0 Hz, C(H)Ar), 5.28 (1H, d, J=9.0 Hz, C(H)Ar), 5.29-5.38 (2H, m, CH₂CH=CH₂), 5.83-5.95 (1H, m, CH₂CH=CH₂), 7.54 (2H, d, J=9.2 Hz, Ar), 8.22 (2H, d, J=9.2 Hz, Ar)

Example 16

(3S,4S)-3-[(R)-1-(tert-Butyldimethylsilyloxy)ethyl]-4-[(RS)-1-(hydroxycarbonyl)ethyl]-1-(p-nitrobenzyloxy-carbonyl)methyl-2-azetidinone

- 5 Triethylamine (32 μ l, 0.233 mmol), formic acid (7.3 μ l, 0.194 mmol), triphenylphosphine (2.0 mg, 0.0077 mmol) and palladium acetate (1.6 mg, 0.0038 mmol) were added to a solution of the compound (48 mg, 0.0775 mmol) as obtained in Example 14 in 1,4-dioxane (1.5 ml), successively, under an argon stream, and the reaction mixture was stirred at 100°C for one hour. The mixture was cooled, admixed with potassium hydrogensulfate (5 ml) and extracted with ethyl acetate (25 ml). The ethyl acetate layer was washed with aqueous saturated sodium chloride solution, dried over sodium sulfate and concentrated under reduced pressure to give 45 mg (quant.) of the subejct compound as a mixture consisting of the α (R-isomer)/ β (S-isomer) at a ratio of 1/1.7.

10 NMR (CDCl_3): δ 0.02-0.08(6H,m,Si(CH₃)₂), 0.85 and 0.86(total 9H,each s,SiC(CH₃)₃), 1.19-1.28(6H,m,CH₃CHOSi and CH₃CHCO₂H), 2.71(0.37H,dd,J=7.1 and 7.4Hz,CHCO₂H), 2.83-2.92(1H,m,CHCO₂H and C3-H), 3.05(0.63H,dd,J=2.0 and 6.0Hz, C3-H), 3.90-4.32 (4H,m,CHOSi,CH₂CO₂ and C4-H), 5.08-5.33 (2H,m,CO₂CH₂), 7.45-7.70(4H,m,Ar)

Example 17

25 (3S,4S)-3-[(R)-1-(tert-Butyldimethylsilyloxy)ethyl]-4-[(RS)-1-cyclohexan-2-on-1-yl]-1-(p-nitrobenzyloxycarbonyl)methyl-2-azetidinone

- 30 Triethylamine (13.6 μ l, 0.0972 mmol), formic acid (2.9 μ l, 0.078 mmol), triphenylphosphine (1.0 mg, 0.0038 mmol) and palladium acetate (0.4 mg, 0.0019 mmol) were added to a solution of the compound (20 mg, 0.0331 mmol) as obtained in Example 15 in tetrahydrofuran (1 ml), successively, under an argon stream, and the reaction mixture was stirred at room temperature for one and a half hours. The mixture was admixed with aqueous saturated sodium chloride solution and extracted with ethyl acetate. The ethyl acetate layer was dried over sodium sulfate and concentrated under reduced pressure to give the subejct compound as a mixture consisting of the α (R-
- 35

isomer)/ β (S-isomer) at a ratio of 1/1.3. Purification of the residue by column chromatography (5 g of silica gel; a 5/1 mixture of hexane/ethyl acetate) yielded 4 mg (23 %) of the α -form and 7 mg (41 %) of the β -form.

5 α -Form (R-isomer): NMR (CDCl_3): δ 0.03 and 0.06 (total 6H, each s, $\text{Si}(\text{CH}_3)_2$), 0.86 (9H, s, $\text{SiC}(\text{CH}_3)_3$), 1.22 (3H, d, $J=5.9\text{Hz}$, CH_3CHOSi), 1.23-1.44 (1H, m), 1.55-1.76 (2H, m), 1.80-1.97 (1H, m), 2.04-2.18 (2H, m), 2.18-2.31 (2H, m), 2.61-2.75 (1H, m, $\text{C}_4\text{CHC=O}$), 2.80 (1H, dd, $J=2.0$ and 5.3Hz , C-3H), 3.88 (1H, dd, $J=2.0$ and 9.9Hz , C4-H), 4.09 (1H, d, $J=17.8\text{Hz}$, CH(H)CO₂PNB), 4.10-4.20 (1H, m, CHOSi), 4.23 (1H, d, $J=17.8\text{Hz}$, CH(H)CO₂PNB), 5.18 (2H, q, $J=13.2\text{Hz}$, CH₂Ar), 7.50 (2H, d, $J=9.2\text{Hz}$, Ar), 8.22 (2H, d, $J=9.2\text{Hz}$, Ar)

10 β -Form (S-isomer): NMR (CDCl_3): δ 0.01 and 0.05 (total 6H, each s, $\text{Si}(\text{CH}_3)_2$), 0.84 (9H, s, $\text{SiC}(\text{CH}_3)_3$), 1.24 (3H, d, $J=5.9\text{Hz}$, CH_3CHOSi), 1.56-1.89 (3H, m), 1.95-2.21 (3H, m), 2.23-2.39 (2H, m), 2.56-2.66 (1H, m, $\text{C}_4\text{CHC=O}$), 2.95 (1H, dd, $J=2.6$ and 7.2Hz , C3-H), 3.90 (1H, d, $J=17.8\text{Hz}$, C(H)HCO₂PNB), 4.18 (1H, d, $J=17.8\text{Hz}$, C(H)HCO₂PNB), 4.09-4.21 (1H, m, CHOSi), 4.28-4.31 (1H, m, C4-H), 5.26 (2H, d, $J=9.2\text{Hz}$, CH₂Ar), 7.55 (2H, d, $J=9.2\text{Hz}$, Ar), 8.23 (2H, d, $J=9.2\text{Hz}$, Ar)

[Industrial Applicability]

As has been described above, the present invention can provide a process of converting the 4-position substituent of the azetidinone derivatives of the general formula (1) in one step to produce the derivatives of the general formula (3), and also can allow the derivatives (3) and (4) to undergo a decarboxylation reaction to give the azetidinone derivatives of the general formula (5), thereby enabling the azetidinone derivatives of the general formula (5) to be synthesized easily in a decreased number of steps as compared with the conventionally known procedures.